

PHYSICIAN ORDER SET :
Iron Sucrose VENOFR

CDH 208-244 – Approved - Page 1 of 3

Patient: _____ DOB: _____ Gender: _____

Patient Phone #: _____ Height: _____ Weight: _____

Diagnosis: _____ ICD-10 Code: _____

Treatment Start Date: _____

Provider Facility Name: _____ Provider Facility Address: _____

Ordering Provider: _____ Date: _____

Signature: _____

Communication Orders

☐ **Provider and Nurse Communication**

Routine, Once, Starting when released.

Total dose of iron sucrose should not exceed 1000mg per treatment course.

Medications

☐ **iron sucrose (VENOFER)**

To be given per site medication administration guidelines.

For doses ≥ 300 mg, observe patients for signs and symptoms of hypersensitivity reaction during and following administration for at least 30 minutes.

Dose (expressed in mg of elemental iron)

☐ 100 mg ☐ 200 mg ☐ 300mg ☐ 400mg ☐ Other _____

Frequency

☐ 2 times a week, at least 2 days apart

☐ Every _____

Total doses to be given:

☐ 3 ☐ 5 ☐ Other _____

PHYSICIAN ORDER SET :
Iron Sucrose VENOFER

CDH 208-244 – Approved - Page 2 of 3

Catheter Management

	Interval	Defer Until	Duration
☐ Line Access Routine, As needed, Starting when released, Until Specified. As needed. Until Specified. Insert peripheral IV, or access peripheral, or central venous access device to provide treatment.	PRN		PRN
☐ alteplase (CATHFLO) 1mg/mL injection 2 mg 2 mg, Intracatheter, As needed, line care, Starting when released. For central venous access device requiring clearance. May repeat once per lumen. For patients greater than 30 kg: 2 mg in 2 mL using a 10 mL syringe. For patient less than or equal to 30 kg: instill a volume equal to 110% of the internal lumen of CVAD, not to exceed 2 mg in 2 mL.	PRN		PRN
☐ lidocaine-prilocaine (LMX 4) cream Topical, As needed, apply prior to the PIV insertion or port access, Starting when released. For TOPICAL Use Only	PRN		PRN
☐ heparin (PF) 100 unit/mL flush 5 mL 5 mL, Intravenous, As needed, line care, Starting when released	PRN		PRN
☐ heparin (PF) 10 unit/mL flush 3 mL 3 mL, Intravenous, As needed, line care, Starting when released	PRN		PRN
☐ heparin (PF) 10 unit/mL flush 5 mL 5 mL, Intravenous, As needed, line care, Starting when released	PRN		PRN
☐ heparin (PF) 1000 unit/mL catheter injection 2 mL 2 mL, Intracatheter, As needed, APHERESIS LINE CARE ONLY. HEPARIN MUST BE WITHDRAWN FROM EACH LUMEN PRIOR TO FLUSHING OR INFUSING THROUGH THE APHERESIS CATHETER, Starting when released	PRN		PRN
☐ sodium chloride (NS) 0.9% syringe flush 3 mL 3 mL, Intravenous, As needed, Line care per institutional guidelines, Starting when released.	PRN		PRN
☐ sodium chloride (NS) 0.9% syringe flush 10 mL 10 mL, Intravenous, As needed, Line care per institutional guidelines, Starting when released.	PRN		PRN
☐ sodium chloride (NS) 0.9% syringe flush 20 mL 20 mL, Intravenous, As needed, Line care per institutional guidelines, Starting when released.	PRN		PRN
☐ sodium chloride 0.9% infusion 20 mL/hr, Intravenous, Continuous PRN, Keep vein open to provide treatment, Starting when released	PRN		PRN
☐ D5W infusion 20 mL/hr, Intravenous, Continuous PRN, Keep vein open to provide treatment, Starting when released, For 24 hours	PRN		PRN

Emergency Medications/Anaphylaxis

	Interval	Defer Until	Duration
☐ Provider and Nurse Communication Routine, Once, Starting when released. Treatment of SEVERE reaction (ANAPHYLAXIS): hypotension, throat swelling, wheezing, respiratory distress, or decreased oxygen saturation. Stop the infusion and treat with epinephrine FIRST. Notify provider and emergency personnel, administer oxygen as needed, monitor vital signs and proceed with administering adjunct HYPERSENSITIVITY medications as clinically indicated.	PRN		PRN

PHYSICIAN ORDER SET :
Iron Sucrose VENOFER

CDH 208-244 – Approved - Page 3 of 3

☐	EPINEPHrine (ADRENALIN) injection 0.3 mg 0.3 mg, Intramuscular, As needed, anaphylaxis, Administer FIRST for anaphylaxis. May repeat times 1 dose, Starting when released For 2 doses. Pharmacy's Suggested Dose Instructions: Epinephrine 1:1000 equivalent to 1 mg/mL	PRN	PRN
☐	sodium chloride 0.9% bolus 1,000 mL 1,000 mL, Intravenous, Once as needed, Hypotension, Starting when released, For 1 dose	PRN	PRN
☐	Oxygen Therapy – Non-Rebreather Routine Select a Mode of Therapy: Non-Rebreather Titrate Oxygen and use the most appropriate device to maintain Target Oxygen saturation during Activity/Titration: Yes Min SpO2 (%): 94	PRN	PRN

Hypersensitivity

	Interval	Defer Until	Duration
☐	PRN		Until discont'd
Provider and Nurse Communication Routine, Until discontinued, Starting when released. Until Specified. Treatment for mild-moderate infusion reaction: Stop the infusion, notify provider and emergency personnel, administer oxygen as needed, monitor vital signs and proceed with administering medications as clinically indicated. If ANAPHYLAXIS reaction, refer to Emergency Medications section.			
☐	PRN		Until discont'd
albuterol 2.5mg/3 mL (0.83%) nebulizer solution 2.5 mg, Nebulization, Once as needed, wheezing, shortness of breath, Starting when released, For 1 dose			
☐	PRN		Until discont'd
acetaminophen (TYLENOL) tablet 975 mg 975 mg, Oral, Once as needed, fever, Starting at treatment start time, For 1 dose			
☐	PRN		Until discont'd
diphenhydramine (BENADRYL) injection 25 mg 25 mg, Intravenous, As needed, itching, hives. Begin with 25 mg. If patient has continued reaction, administer 25 mg, Starting when released, For 1 dose			
☐	PRN		Until discont'd
methylprednisolone sodium succinate (PF) (SOLU-Medrol) injection 40 mg 40 mg, Intravenous, Once as needed, Adjunct treatment for mild-moderate, or SEVERE reaction, Starting when released, For 1 dose			
☐	PRN		Until discont'd
famotidine (PEPCID) injection 20 mg 20 mg, Intravenous, Once as needed, Adjunct treatment for mild-moderate, or SEVERE reaction, Starting when released, For 1 dose To be administered along with H1 antihistamine and famotidine			
☐	PRN		Until discont'd
ondansetron (PF) (ZOFran) injection 4 mg 4 mg, Intravenous, As needed, nausea, vomiting, may repeat x 1 dose, Starting when released, For 2 doses			
☐	PRN		Until discont'd
cetirizine (Zyrtec) tablet 10 mg 10 mg, Oral, Once as needed, Adjunct treatment for mild-moderate, or SEVERE reaction, Starting at treatment start time, For 1 dose. If patient unable to tolerate cetirizine, administer fexofenadine if available. HOLD IF: given fexofenadine.			
☐	PRN		Until discont'd
fexofenadine (Allegra) tablet 180 mg 180 mg, Oral, Once as needed, Adjunct treatment for mild-moderate, or SEVERE reaction, Starting at treatment start time, For 1 dose. Administer only if unable to tolerate cetirizine. HOLD IF: given cetirizine.			